



0-0-8

GUARANTEED ANALYSIS

Soluble Potash (K₂O) 8%

DERIVED FROM: Muriate of Potash

ALSO CONTAINS NON-PLANT FOOD INGREDIENTS:

0.15% Humic Acid (derived from humic shale)

0.06% Kelp (microbe food)

This material is not considered hazardous by the OSHA Hazard Communication Standard (29 CFR 1910.1200)

NOTE: When mixing EMPAC F with other chemicals, all chemical labels should be read in their entirety to avoid potential and undesirable reactions, ensuring the compatibility based on the pH.

STORAGE: No special handling required. Do not store above 90°F. Do not freeze. Keep out of direct sunlight for prolonged periods of time.

SAFETY: Do not take internally. Avoid eye or prolonged skin contact. Should contact occur, wash with water.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.html>

NET CONTENTS:

275 GAL (1,041 Liters) | 2612.50 lbs (1185.01kgs)

DENSITY: 9.5 lbs/gallon at 68°F

NOTICE: Verdeflo LLC. Warrants this product to the chemical description and for it's specific intended use only. This warrantee is only valid when the product is used in accordance with the directions also described on this label. It is understood that it is impossible to eliminate ALL risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may be a result of many factors, such as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Verdeflo LLC. In no case shall Verdeflo LLC. be held liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. Verdeflo LLC. makes no warranties in any other express or implied warranty except as stated above.



7500 W Lake Mead Blvd, Suite 9 #127 • Las Vegas NV, 89128
702-591-6838 • FAX: 702-549-2175

DIRECTIONS FOR USE

BERRIES (Blackberries, Blueberries, Raspberries, Strawberries)

- Pre-bloom: Apply 1-3 gallons of EMPAC F per acre diluted with 15 gallons of water.
- Fruit Set: Apply 1-3 gallons of EMPAC F per acre diluted with 15 gallons of water.

GRAPES (Table & Wine)

- Pre-Bloom: Apply 1-5 gallons of EMPAC F per acre through drip irrigation.
- Fruit Set: Apply 1-3 gallons of EMPAC F per acre through drip irrigation.

PINTO BEANS/RED BEANS

- Flowering: Apply 1-3 gallons of EMPAC F per acre diluted with 15 gallons of water (foliar or shanked in).

POTATOES

- Tuber Initiation: Apply 1-3 gallons of EMPAC F per acre diluted with 15 gallons of water (foliar or shanked in).

SOYBEANS

- Emergence or with 1st glyphosate: Apply 1-3 gallons of EMPAC F per acre diluted with 15 gallons of water.

TOMATO/PEPPERS

- Pre-bloom: Apply 1-3 gallons of EMPAC F per acre with a minimum of 15-20 gallons of water.
- Fruit Set: Apply 1-3 gallons of EMPAC F per acre with a minimum of 15-20 gallons of water.

TREES (Fruit & Nut)

- Pre-Bloom: Apply 1-4 gallons of EMPAC F per acre through irrigation. (Root Drench).
- Fruit Set: Apply 1-4 gallons of EMPAC F per acre through irrigation. (Root Drench).

VEGETABLE (Row Crops) Broccoli, Cauliflower, Cabbage, Celery, Head Lettuce, Mix Lettuce, Spring Mix, Spinach

- Thinning: Apply 1-3 gallons of EMPAC F per acre to 1st Side Dress Application. (Can be applied through water runs also).

LOT NUMBER: _____

KEEP OUT OF REACH OF CHILDREN